

# Referenser i Ortopediskt Magasin nr 4, 2020 som inte fick plats i tidningen

## Omvårdnad vid geriatrisk traumatologi Sidan 28-29 i OM nr 4 2020

### Referenser:

1. Asplund, K., Gustafson, Y., Jacobsson, C., Bucht, G., Wahlin, A., Peterson, J., Blom, J. O. & Angquist, K. A. 2000. Geriatric-based versus general wards for older acute medical patients: a randomized comparison of outcomes and use of resources. *J Am Geriatr Soc*, 48, 1381-8.
2. Lundström, M., Olofsson, B., Stenvall, M., Karlsson, S., Nyberg, L., Englund, U., Borssen, B., Svensson, O. & Gustafson, Y. (2007). Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clinical and Experimental Research*, 19(3), 178-186.
3. Vårdhandboken Trycksår Hämtad 2020-11-28 <https://www.varldhandboken.se/varld-och-behandling/hud-och-sar/trycksar/oversikt/>
4. Vårdhandboken Munhälsa Hämtad 2020-11-28 <https://www.varldhandboken.se/varld-och-behandling/basal-och-preventiv-omvardnad/munhalsa/>
5. Vårdhandboken Fallprevention Hämtad 2020-11-20 <https://www.varldhandboken.se/varld-och-behandling/basal-och-preventiv-omvardnad/fallprevention/riskbedomning/>
6. <https://www.hifeprogram.se/>
7. Edwards, B.A., O'Driscoll, D.M., Ali, A., Jordan, A S., Trinder, J. & Malhotra, A. (2010). Aging and sleep: physiology and pathophysiology. *Seminars in Respiratory and Critical Care Medicine*, 31(5), 618-633. Brooke, R. C. & Griffiths, C. E. Aging of the Skin. I: Raymond C. Tallis H. M. F. 2003, Ed *Brocklehurst's textbook of Geriatric Medicine and Gerontology*. 6th Edition: London: Churchill Livingstone, sid 1269-1275. Clarke, B. L. & Khosla, S. 2010. Physiology of bone loss. *Radiol Clin North Am*, 48, 483-95. Socialstyrelsen Jämlik hälsa, vård och omsorg. Hämtad 2020-11-20. <https://www.socialstyrelsen.se/utveckla-verksamhet/jamlik-halsa-varld-och-omsorg/> Stenvall, M., Olofsson, B., Lundström, M., Englund, U., Borssen, B., Svensson, O., Nyberg, L. & Gustafson, Y. (2007). A multidisciplinary, multifactorial intervention program reduces postoperative falls and injuries after femoral neck fracture. *Osteoporosis International*, 18(2), 167-175. Svensk sjuksköterskförening, Svenska Läkarsällskapet och Dietisternas Förbund 2019. Personcentrerad vård – en kärnkompetens för god och säker vård. Hämtad: 2020-11-28 <https://swenurse.se/download/18.9f73344170c003062310d6/1583937715986/personcentrera-d%20v%C3%A5rd%202019.pdf>

# Geriatriskt trauma – om att ramla i livets utförsbacke

## Sidan 33-35 i OM nr 4 2020

### Referenser

1. McLachlan AJ, Bath S, Naganathan V, Hilmer SN, Le Couteur DG, Gibson SJ, et al. Clinical pharmacology of analgesic medicines in older people: impact of frailty and cognitive impairment. *Br J Clin Pharmacol*. 2011;71(3):351-64.
2. Edelstein DM, Aharonoff GB, Karp A, Capla EL, Zuckerman JD, Koval KJ. Effect of postoperative delirium on outcome after hip fracture. *Clin Orthop Relat Res*. 2004(422):195-200.
3. Griffiths R, Babu S, Dixon P, Freeman N, Hurford D, Kelleher E, et al. Guideline for the management of hip fractures 2020. *Anaesthesia*.n/a(n/a).
4. A WE, Ekerstad N, Alfredsson J, Johanson M, Metzner C, Wilhelmson K, et al. [Frailty]. *Lakartidningen*. 2020;117.
5. Ritcey B, Pageau P, Woo MY, Perry JJ. Regional Nerve Blocks For Hip and Femoral Neck Fractures in the Emergency Department: A Systematic Review. *Cjem*. 2016;18(1):37-47.
6. Olofsson C, Nygård EB, Bjersten AB, Hessling A. Low-dose bupivacaine with sufentanil prevents hypotension after spinal anesthesia for hip repair in elderly patients. *Acta Anaesthesiol Scand*. 2004;48(10):1240-4.
7. Heffernan DS, Thakkar RK, Monaghan SF, Ravindran R, Adams CA, Jr., Kozloff MS, et al. Normal presenting vital signs are unreliable in geriatric blunt trauma victims. *J Trauma*. 2010;69(4):813-20.
8. Salottolo KM, Mains CW, Offner PJ, Bourg PW, Bar-Or D. A retrospective analysis of geriatric trauma patients: venous lactate is a better predictor of mortality than traditional vital signs. *Scand J Trauma Resusc Emerg Med*. 2013;21:7.
9. White SM, Moppett IK, Griffiths R, Johansen A, Wakeman R, Boulton C, et al. Secondary analysis of outcomes after 11,085 hip fracture operations from the prospective UK Anaesthesia Sprint Audit of Practice (ASAP-2). *Anaesthesia*. 2016;71(5):506-14.
10. Sessler DI, Bloomstone JA, Aronson S, Berry C, Gan TJ, Kellum JA, et al. Perioperative Quality Initiative consensus statement on intraoperative blood pressure, risk and outcomes for elective surgery. *Br J Anaesth*. 2019;122(5):563-74.
11. Cedborg AI, Sundman E, Boden K, Hedstrom HW, Kuylenstierna R, Ekberg O, et al. Pharyngeal function and breathing pattern during partial neuromuscular block in the elderly: effects on airway protection. *Anesthesiology*. 2014;120(2):312-25.
12. Klestil T, Röder C, Stotter C, Winkler B, Nehrer S, Lutz M, et al. Impact of timing of surgery in elderly hip fracture patients: a systematic review and meta-analysis. *Sci Rep*. 2018;8(1):13933.
13. Accelerated surgery versus standard care in hip fracture (HIP ATTACK): an international, randomised, controlled trial. *Lancet*. 2020;395(10225):698-708.

# Operation för höftfraktur inom 24 timmar – är det dags att tänka nytt?

## Sidan 46-47 i OM nr 4 2020

### Referenser

1. Orosz GM, Magaziner J, Hannan EL. Association of timing of surgery for hip fracture and patient outcomes. JAMA. 2004; 291(14): 1738-1743
2. Simunovic N, Devereaux PJ, Sprague S, et al. Effect of early surgery after hip fracture on mortality and complications: systematic review and meta-analysis. CMAJ. 2010; 182(15): 1609-1616
3. Målnivåer för rörelseorganens sjukdomar. Socialstyrelsen.  
<https://www.socialstyrelsen.se/globalassets/sharepoint-dokument/artikelkatalog/nationella-riktlinjer/2020-9-6906.pdf>
4. Rikshöft. Årsrapport 2020. [https://04e8d8b0-c67b-4aa0-a7e7-d272a37c2285.filesusr.com/ugd/3ac01b\\_7ed6560d1255425b964d6b19833f148e.pdf](https://04e8d8b0-c67b-4aa0-a7e7-d272a37c2285.filesusr.com/ugd/3ac01b_7ed6560d1255425b964d6b19833f148e.pdf)
5. Turesson E. A 30-year journey in hip fracture care. An evaluation of how care process development affect lead-times and outcome. Lunds universitet. 2019. Akademisk avhandling. ISBN 978-91-7619-764-6
6. Eriksson M, Kelly-Pettersson P, Stark A, et al. 'Straight to bed' for hip-fracture patients: A prospective observational cohort study of two fast-track systems in 415 hips. Injury, International Journal of the Care of the Injured; 43 (2012) 2126-2131
7. Leer-Salvesen S, Engesæter LB, Dybvik E, et al. Does time from fracture to surgery affect mortality and intraoperative medical complications for hip fracture patients? An observational study of 73 557 patients reported to the Norwegian hip fracture register. J Bone Joint Surg. 2019; 101-B: 1129-1137
8. Greve K, Modig K, Talback M, et al. No association between waiting time to surgery and mortality for healthier patients with hip fracture: a nationwide Swedish cohort of 59,675 patients. Acta Orthop. 2020; 91: 396-400
9. Borges FK, Bhandari M, Guerra-Farfan E, et al. Accelerated surgery versus standard care in hip fracture (HIP ATTACK): an international, randomised, controlled trial. The Lancet. 2020; 395: 698-708
10. Pincus D, Ravi B, Wasserstein D, et al. Association Between Wait Time and 30-Day Mortality in Adults Undergoing Hip Fracture Surgery. JAMA. 2017; 318: 1994-2003
11. Okada E, Inukai K, Aoyama H. Wait Time for Hip Fracture Surgery and Mortality. JAMA. 2018; 319: 2233
12. Sheehan KJ, Sobolev B, Guy P. Mortality by timing of hip fracture surgery. Factors and relationships at play. J Bone Joint Surg. 2017; 99: e106(1-9)
13. Reid IR. A broader strategy for osteoporosis interventions. Nature Reviews Endocrinology. 2020; 16: 333-339. DOI: 10.1038/s41574-020-0339-7
14. Ekdahl AW, Ekerstad N, Alfredsson J, et al. Skörhetsbegreppet viktigt för att förstå den äldre patientens behov. Läkartidningen. 2020;117:F3HE
15. Aggarwal A, Harris IA, Naylor JM. Patient preferences for emergency or planned hip fracture surgery: a cross-sectional study. J. Orthop. Surg. Res. 2016; 11: 120

16. Björkman-Björkelund, K. (2008) Acute confusional state in elderly patients with hipfracture. Identification of risk factors and intervention using a prehospital and perioperative management program. Lund University
17. Rogmark C. Further refinement of surgery will not necessarily improve outcome after hip fracture. *Acta Orthop.* 2020; 91(2): 123-124
18. Neuerburg C, Förch S, Gleich J, et al. Improved outcome in hip fracture patients in the aging population following co-managed care compared to conventional surgical treatment: a retrospective, dual-center cohort study. *BMC Geriatr* 19, 330 (2019).  
<https://doi.org/10.1186/s12877-019-1289-6>
19. Svenska knäprotesregistret Årsrapport 2020. [http://myknee.se/pdf/SVK\\_2020\\_v1.0.pdf](http://myknee.se/pdf/SVK_2020_v1.0.pdf)
20. Symons N, Almoudaris AM, Nagpal K, et al. An observational study of the frequency, severity, and etiology of failures in postoperative care after major elective general surgery. *Ann Surg.* 2013 Jan;257(1):1-5. doi: 10.1097/SLA.0b013e31826d859b
21. Exercise for preventing falls in older people living in the community: Update of Cochrane Systematic Review. [Apr 27]. [https://www.who.int/docs/default-source/physical-activity/call-for-consultation/evidence-on-falls-prevention.pdf?sfvrsn=2708e33\\_4](https://www.who.int/docs/default-source/physical-activity/call-for-consultation/evidence-on-falls-prevention.pdf?sfvrsn=2708e33_4)
22. Meehan AJ, Maher AB, Brent L, et al. The International Collaboration of Orthopaedic Nursing (ICON): Best practice nursing care standards for older adults with fragility hip fracture. *Int J Orthop Trauma Nurs.* 2019; 32: 3-26

## **Replik på: Operation för höftfraktur inom 24 timmar – är det dags att tänka nytt?**

### **Sidan 48-49 i OM nr 4 2020**

#### **Referenser**

1. Tomioka, S., Rosenberg, M., Fushimi, K., & Matsuda, S. (2020). An analysis of equity in treatment of hip fractures for older patients with dementia in acute care hospitals: observational study using nationwide hospital claims data in Japan. *BMC Health Services Research*, 20(1), 1-12.
2. Membership of the Working Party, Griffiths, R., Alper, J., Beckingsale, A., Goldhill, D., Heyburn, G., ... & White, S. (2012). Management of proximal femoral fractures 2011: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia*, 67(1), 85-98.
3. Shiga, T., Wajima, Z. I., & Ohe, Y. (2008). Is operative delay associated with increased mortality of hip fracture patients? Systematic review, meta-analysis, and meta-regression. *Canadian Journal of Anesthesia*, 55(3), 146.
4. Uzoigwe, C. E., Burnand, H. G. F., Cheesman, C. L., Aghedo, D. O., Faizi, M., & Middleton, R. G. (2013). Early and ultra-early surgery in hip fracture patients improves survival. *Injury*, 44(6), 726-729.
5. Bottle, A., & Aylin, P. (2006). Mortality associated with delay in operation after hip fracture: observational study. *Bmj*, 332(7547), 947-951.

6. Pincus, D., Ravi, B., Wasserstein, D., Huang, A., Paterson, J. M., Nathens, A. B., ... & Wodchis, W. P. (2017). Association between wait time and 30-day mortality in adults undergoing hip fracture surgery. *Jama*, 318(20), 1994-2003.
7. Borges, F. K., Bhandari, M., Guerra-Farfan, E., Patel, A., Sigamani, A., Umer, M., ... & Teixidor-Serra, J. (2020). Accelerated surgery versus standard care in hip fracture (HIP ATTACK): an international, randomised, controlled trial. *The Lancet*, 395(10225), 698-708.
8. Greve K, Modig K, Talback M, et al. No association between waiting time to surgery and mortality for healthier patients with hip fracture: a nationwide Swedish cohort of 59,675 patients. *Acta Orthop*. 2020; 91: 396-400
9. Aggarwal A, Harris IA, Naylor JM. Patient preferences for emergency or 189 planned hip fracture surgery: a cross-sectional study. *J. Orthop. Surg. Res.* 190 2016; 11: 120
10. Vestergaard, P., Rejnmark, L., & Mosekilde, L. (2007). Has mortality after a hip fracture increased?. *Journal of the American Geriatrics Society*, 55(11), 1720-1726.
11. Pedersen, A. B., Ehrenstein, V., Szépligeti, S. K., Lunde, A., Lagerros, Y. T., Westerlund, A., ... & Sørensen, H. T. (2017). Thirty-five-year trends in first-time hospitalization for hip fracture, 1-year mortality, and the prognostic impact of comorbidity: a Danish nationwide cohort study, 1980–2014. *Epidemiology*, 28(6), 898-905.
12. Gundel, O., Thygesen, L. C., Gögenur, I., & Ekeloef, S. (2020). Postoperative mortality after a hip fracture over a 15-year period in Denmark: a national register study. *Acta orthopaedica*, 91(1), 58-62.